



INTACT

INDIGENOUS PEOPLES TRANSFORMING
ALZHEIMER'S CARE TRAINING

Part One: Making a Diagnosis of Cognitive Impairment



DISCLOSURE

Barak Gaster, MD

No financial relationships with any commercial company which is relevant to this presentation to disclose.

I have no relevant financial relationships with any companies whose primary business is producing, marketing, selling, re-selling, or distributing any healthcare products or services which are used by or on patients.

Three Part Education Series

Part #1

Evaluate cognition

Part #2

Set plan for newly
diagnosed patient

Part #3

Tips for managing
dementia

What makes dementia hard?


- The diagnosis in early stages can be tricky.
- Tempting for patients and us to skip over it.
- But we can intervene meaningfully if we identify it earlier!

Diagnosis: a Path to Better Care



- Better communication. Better support. Improved involvement of family.
- Care for patient actually gets easier, less chaotic.
- Steps you can take now which can improve cognition.

Dementia Shock Ahead



Numbers will double
in next 20 years

6.9 million Americans now

35% of all people over age 85

Changes for American Indians

- American Indians (AIs) are living longer! Which is good news. Life expectancy for AIs has increased by 68% in the past 50 years.
- The number of American Indians over age 65 is expected to double in coming years.
- The number over 85 is expected to increase by 7-fold between now and 2050.

Challenge Ahead

- Having more older tribal members is a welcome thing. Immense value in their wisdom, experience, and contributions to American Indian communities.
- But huge challenge also: facing a skyrocketing number of American Indians with dementia.
- Coming problems for families, health clinics, tribal leadership, and community supports.

Learning Objectives

- Perform a cognitive evaluation.
- Identify reversible and contributing causes of cognitive impairment you can fix.
- Distinguish “Mild Cognitive Impairment” from Dementia



72 year old woman in your office for a wellness visit. Generally very healthy.

Hypertension, osteoarthritis.

Medications: Losartan, rare ibuprofen, rare zolpidem.

She happens to mention, “I’ve been worried about my memory.”

Should you:

- X.** Reassure her? You might say “Don’t worry, it’s so common and normal to get more forgetful with age.”
- X.** Automatically refer to a specialist ?
- X.** Change course, try to do a cognitive assessment then and there (and then run 35 minutes late?)

D. Offer she schedule an efficient, follow-up visit with you, so you can evaluate this further.



~ BE AWARE



- Patient mentions a concern
- Family member raises a concern
- Annual wellness visit questionnaire?
- You or other clinic staff notice changes



72 year old woman in your office for a wellness visit. Generally very healthy.

Hypertension, osteoarthritis.

Medications: Losartan, rare ibuprofen, rare zolpidem.

She happens to mention, “I’ve been worried about my memory.”

You need a little more information

How strongly should you say: “I’m worried. It’s so important we get you a full evaluation.”

(sometimes concerns can be chalked up easily to normal aging)

Or are there worrisome signs? → Clues help to decide ...

Clues to help you decide

Normal: Misplacing keys. Forgetting why I came up the stairs.

Slower to remember a word or a name, but later it comes to you.



WARNING SIGNS

Clues to help you decide

Versus serious signs:

1. Forgetting things that just happened.

Such as: Repeating same question 30 minutes later.

2. Harder to do complex task that used to be easy.

Such as: Trouble making a complex recipe or organizing documents.

3. Suddenly unsure where you are in place you've been often.

Example: Suddenly disoriented in a building you should know well.

Decide:

Anyone with a concern: it is perfectly fine to offer a cognitive evaluation.

Framework for how strongly to insist

(sometimes you can reassure)

But fine to always still offer an evaluation.

Normal: Misplacing keys. Why did I come up stairs? Or slower to remember a word or remember a name, but later it comes to you.

Be aware of far more worrisome signs:

1. Forgetting things that just happened. (repeating questions)
2. Harder to do complex task that used to be easy.
3. Suddenly unsure where you are in place you've been often.

Decision Aid

One-pager to use with patients is available in a folder at your clinic.

Quick Questions if Memory is a Concern

1. Have you noticed that you forget things that just happened more often?
For example: Repeating the same question or the same story 30 minutes later.

YES	NO	UNSURE
-----	----	--------

2. Have you noticed it's difficult to finish a complex task that used to be easy for you?
For example: Cooking a complex recipe, organizing your documents, or putting up outdoor holiday lights.

YES	NO	UNSURE
-----	----	--------

3. Have you noticed being unsure where you are in a place you've been to many times?
For example: Becoming disoriented on a usual route or in a building you know well.

YES	NO	UNSURE
-----	----	--------



in for an annual
wellness visit.


She happens to
mention, “I’ve been
worried about my
memory.”

“I’m really having
trouble keeping
things straight. It’s
really not right.”

You decide yes

Schedule a Cognitive Evaluation Visit

- Strongly encourage a **family member** or a close friend to come to that next visit.



Three Parts to a Dedicated Visit to Evaluate Cognitive Concerns

1. **Cognitive Checklist:** reversible causes and other factors you can fix.
2. **Assess cognitive function** e.g. with a MoCA.
3. **Get input** from a family member (e.g. with AD8).

Cognitive Checklist

- ☐ **Labs** B12 + thyroid
- ☐ **Med list** Sedating/ anticholinergic meds
 Oxybutynin, Tylenol PM, benzodiazepines, Ambien
- ☐ **Alcohol** Even mild-to-moderate drinking can
 impair cognition once people > age 70
- ☐ **Conditions:** Sleep apnea, hearing loss, depression

Checklist now available in your EHR

HPI - Text Macros

Evaluating Cognitive Function

The assessment is a combination of...

MoCA test


+

Family input

MoCA Montreal Cognitive Assessment

The best validated tool for identifying mild cognitive impairment in the primary care setting.

More sensitive than MMSE.



Family / Observer Input

History questions to ask family member or a close friend:

- Are they repeating the same question 30 min later?
- Losing ability to do complex tasks that they once found easy to do? (e.g. complex recipe or outdoor lighting)
- Getting disoriented in a familiar place?

Remember these three questions, or....

Use 8-item interview guide: **the AD8** now in folders at your clinics

Observer Input

AD8

validated tool to
get input from a
family member
or close friend.

Remember, "Yes, a change" indicates that there has been a change in the last several years caused by cognitive (thinking and memory) problems.	YES, A change	NO, No change
1. Problems with judgment (e.g., problems making decisions, bad financial decisions, problems with thinking)		
2. Less interest in hobbies/activities		
3. Repeats the same things over and over (questions, stories, or statements)		

Delirium vs. Dementia

- MoCA not accurate during an acute illness.
- Transient drop in cognition common in the elderly. That's delirium, not dementia.
- Delirium post hospitalization can last weeks.
- However, delirium is a warning sign →
Odds of dementia (either present now, or developing later) are much higher.

Putting It Together

MoCA result

+

Family input

MoCA less than 20 (out of 30)

+ Observer notes worrisome changes, and

+ Difficulty with *activities of daily living*

= Dementia

Putting It Together

MoCA result

+

Family input

MoCA intermediate 22-26 (out of 30)

+ Observer notes worrisome changes

But still able to dress and cook and clean house

= **Mild Cognitive Impairment**



Normal aging ····· Mild cognitive impairment ····· Dementia


Slower remembering names. Why did I come up the stairs?



Asks the same question 30 minutes later. Can't complete complex task.



Loss of ability to do ADLs: e.g. cooking, driving, or dressing.



Mild Cognitive Impairment (MCI)


- MoCA usually 20-26 plus worrisome changes noted
But activities daily-living **intact**. This is a diagnosis of MCI
(dementia: requires loss independence with activities)
- MCI is a diagnosis in-itself, not simply “early dementia”.
In fact: 30% with MCI **don't** progress to dementia.
But: 70% do progress. Most people who have MCI do
have early Alzheimer's disease. (but not all)
- **MCI Actions!** Focus on brain health: reduce alcohol,
wear hearing aids, sleep apnea.

If the MoCA plus the input from an observer is borderline

- Refer for more testing:
- Specialty referral (neurology or geriatrics), or
- Repeat evaluation in 6-12 months

Either way: address Brain Health Checklist

(Checklist now available in your EHR **HPI - Text Macros**)



Brain Health Checklist

- **Alcohol (and drugs):** Limiting to 0-1 drinks will help.
- **Med List:** Stop sedating or anticholinergics (ask re: OTC's)
- **Contributing Conditions:** Sleep apnea, hearing loss.
- **Exercise:** Daily brisk walks with a friend.
- **Cognitive Stimulation** Encourage social engagement.

Follow-up: set next steps

- **If evaluation looks OK**, discuss ways to keep brain healthy, consider repeat evaluation in 12 months.
- **If you found contributing factors:** counsel on reducing alcohol, wear hearing aids more, treat sleep apnea, treat depression.
- **If looks like mild cognitive impairment or dementia:**
say: “This is a lot to take in and work through. Let’s
Schedule *another appointment* soon to review
what this means and make a plan.”

Make that appointment Encourage family to attend!

Set that plan. Don’t diagnose and adios.

Takeaways

- We can do cognitive evaluations in primary care. Consider using a checklist.
- Having family input (whenever possible) is essential for this evaluation.
- Combine the MoCA and the family input to make your assessment. Consider the AD8.
- Address brain health and key modifiable causes of mild cognitive impairment: Alcohol, sleep apnea, hearing loss, and medications.

Next session

- What are the types of dementia.
- Words to use in discussing the diagnosis.
- Talking about prognosis.
- When to refer to Neurology.
- High value community resources.

Three Part Series

Part #1

Evaluate cognition

Part #2

Set plan for newly
diagnosed patient

Part #3

Tips for managing
dementia



INTACT

INDIGENOUS PEOPLES TRANSFORMING
ALZHEIMER'S CARE TRAINING

Questions and Answers

