



## Part Three: Manage Dementia as it Progresses



# Disclosures

None of the faculty for this educational activity have relevant financial relationships with ineligible companies to disclose.

# Three Part Education Series

Part #1

Make a diagnosis of  
cognitive impairment

Part #2

Set a plan for a newly  
diagnosed patient

**Part #3**

**Manage dementia as  
it progresses**

# Recap of Part #2

Part #1

Make a diagnosis of cognitive impairment

**Part #2**

Set a plan for a newly diagnosed patient

Part #3

Manage dementia as it progresses

## Recap of Part #2 – Set a Plan

- Even if diagnosis is mild cognitive impairment (MCI) mention your high concern for Alzheimer's disease.
- Good to use the MCI diagnosis code and put this in problem list ----> that is better communication/ care.
- Include prognostic uncertainty, include optimism. With support, people can live well. Family should come to all future appointments.
- If visual hallucinations, then refer to a specialist sooner (possible Lewy Body Disease).

# **Learning Objectives Today**


## **Managing Cognitive Impairment**

- Medications to treat cognitive decline.
- Managing agitation and other behavioral symptoms of dementia.
- Communication tips for talking to someone with dementia.
- Advance care planning: for dementia.

# Optimizing Brain Health

- ❑ **Alcohol (and drugs):** Limiting 0-1 drinks will help.
- ❑ **Medications:** Sedating and anticholinergic
- ❑ **Contributing Conditions:** Sleep apnea, hearing loss.
- ❑ **Exercise:** Daily brisk walks with a friend.
- ❑ **Cognitive Stimulation** Social engagement!

# Booklet to Give to Newly Diagnosed



You may be wondering...

- Are there any medications, treatments or lifestyle changes that could help my loved one's memory and thinking?

Early-Stage  
Dementia

Mid-Stage  
Dementia

Late-Stage  
Dementia





Now a return visit one year later.

Repeat MoCA is 21, down from 23.

Daughter says she's still independent with ADLs. So diagnosis is still mild cognitive impairment.

Her daughter asks: what about this new medication I heard about for Alzheimer's disease?

# Lecanemab

- New monoclonal antibody against amyloid plaque.
- RCT's show that it may slow progression of mild cognitive impairment by a few months.
- Twice monthly IV infusion. Major potential harm: significant risk of brain edema/ bleeding.
- If on apixaban or warfarin: not eligible (due to hemorrhage)
- Also to be eligible: patients need a PET-scan or LP with amyloid present. Also need DNA apo-E testing.

# Donepezil and Memantine

- They do not work for early-stage memory loss (MCI).
- It is reasonable to offer a trial of medications. They are safe and easy to prescribe.
- May result in very **small** improvements in cognition and function.
- Symptomatic therapies, not neuroprotective.
- Do not change trajectory of disease.

# Behavioral and Psychological Symptoms of Dementia (BPSD)


- The most troubling part of dementia.
- Agitation, refusal of care, paranoia, yelling, possibly hitting.
- Up to 80% of patients with dementia develop BPSD symptoms at some point in their illness.
- Benzodiazepines: Often make symptoms worse. Use extreme caution.

# BPSD Non-Pharmacological Approaches

- Set reasonable goal of reduction—not elimination—of behaviors.
- Recommend the book, “The 36-Hour Day” (guide for caregivers).
- Consider using a handout with “Tips for talking to a loved one with dementia.”

# Non-Pharmacological Approaches

- Behavioral strategies really do help.
  - Book for caregivers: “The 36-Hour Day.”
  - Tips for communication...



# Talking to someone with dementia

1. Meet them in the reality where they are. Avoid re-grounding. Instead: change subject, change to a related topic.
2. Keep talking directly to someone with dementia (not about them, as if they weren't there.) They might not understand, but that will avoid distress.
3. If aggressive behaviors occur, stay calm. Withdraw if necessary. The moment often passes.

Handout for families, covering this approach, is available at:

[Cognition-PrimaryCare.org](http://Cognition-PrimaryCare.org)

## Antipsychotics: only if extremely serious severe symptoms

- Only if symptoms serious, very severe. Biggest concern is they cause unnecessary, extreme sedation.
- Start very low: Such as **quetiapine** 12.5 mg or **risperidone** 0.25 mg (once daily every evening.)
- Titrate if needed every 3 days. Avoid “as needed” use. (They don’t work fast enough, takes 2 hours to work.)
- Main pitfall: leaving them on too long. (Behaviors eventually improve.) Schedule taper trial in 2 months.
- Warning: Lewy body disease = severe Parkinsonism

# Other medications for PTSD

- Such as citalopram, trazodone, melatonin.
- Limited data they work.
- They can help, but often not much noticeable effect. (They also can also cause side effects.)
- But worth a try, when situation is not extreme, but symptoms are persistent despite behavioral approaches.

# Powerful Tools for Caregivers

Free 6-week classes.

“Take care of yourself while  
you take care of a loved one.”



[www.powerfultoolsforcaregivers.org](http://www.powerfultoolsforcaregivers.org)

# Advance Care Planning for Dementia

Path to better care.




# Dementia is Complex

- People with early dementia may have many years with a good quality of life.
- Often a slow decrease in quality of life: from mild, to moderate, to severe stages.
- It makes sense that people would want different goals for their medical care, from one stage to another.

# **Dementia-specific Advance Directive**

- Developed with input from experts in palliative care, neurology, and geriatrics.
- Tested and refined in primary care.
- Available for anyone to download from:

**[Dementia-directive.org](https://dementia-directive.org)**



*A Simple Way to Document the Medical Care*

*You Would Want If You Had Dementia*

DOWNLOAD THE DEMENTIA DIRECTIVE FORM

INSTRUCTIONS

FAQS

RESOURCES

IN THE NEWS

An advance directive for dementia as featured in the [New York Times](#).

DOWNLOAD THE DIRECTIVE

# Dementia-directive.org

- Brief descriptions of mild, moderate, and severe dementia.
- Below each stage, ability to choose a goals of care option for that stage:

Full code

DNR/DNI

Comfort-focus

## **Best time to offer a Dementia Directive**

- ✓ Before signs of dementia occur.
- ✓ Consider: for everyone over age 65

[www.dementia-directive.org](http://www.dementia-directive.org)

# In Early Impairment: Proxies (DPOA-HC)



- Early-on in the disease, it is important to designate, in a **legal form** who someone want their proxy decision makers to be.
- With alternates.
- Because over 10-15 years, their default (usually their spouse) may no longer be available to serve as their decision maker.

# Tips For Goals of Care Discussions

“Imagine if your loved one could look on themselves now, what might they say they’d want?”

- It is OK to make a recommendation, such as:  
  
“I would recommend not doing CPR. Even if you survived cardiac arrest, your brain is more vulnerable, so you’d be in a worse state afterwards than now. And a stay in the ICU would likely be a hard and very painful time.”



76 year old woman, diagnosis is now mild cognitive impairment.

At serial visits, you cover:

Brain health: reducing alcohol, wearing hearing aids, treating sleep apnea, exercising and remaining socially active.

Help her fill out DPOA form (who would she want as backup if daughter not able?)

Now you're providing better care!

# Take Homes – from Today

- Focus on non-pharmacological approaches to treat behavioral symptoms.
- Antipsychotics can cause severe unnecessary sedation. Only use in very severe situations.
- Advance care planning early in disease: set DPOA/ proxy so that patient can identify an alternate to spouse while they can.

# Take Homes – from the Series

- Early diagnosis is a path to better care.  
Specific actions we can take.
- Use visit dedicated to cognitive evaluation.  
Checklist + MoCA + Input from observer.
- If MCI: Remember ~ 30% don't progress.  
Yet, most do have Alzheimer's. Be honest,  
also provide some reassurance.



## Next session

- Putting it all together.
- Review of the materials in place.
- Value of making the diagnosis.

# CME Information: AMA PRA Category 1 Credit™

Follow the process below to claim CME credit and/or to provide feedback on your learning experience.

## STEP 1

Use your phone to scan QR code:



Or enter URL in browser:  
<https://tinyurl.com/INTACT0912>

## STEP 2

Sign in to WSU  
CloudCME or create  
a new profile

**Submit Activity ID#:**

**3739**

This is in the “Claim  
Credit” area of MyCME

## STEP 3

Complete Evaluation  
and Download CME  
Certificate

This is in the “Evaluations  
& Certificates” area of  
MyCME

For help contact:  
[medicine.cme@wsu.edu](mailto:medicine.cme@wsu.edu)



This activity has been planned and implemented in accordance with the accreditation requirements and policies of the Accreditation Council for Continuing Medical Education (ACCME). The Washington State University (WSU) Elson S. Floyd College of Medicine is accredited by the ACCME to provide continuing medical education for physicians. The WSU Elson S. Floyd College of Medicine designates this activity for a maximum of 2.25 *AMA PRA Category 1 Credits™*. Participants should claim only the credit commensurate with the extent of their participation in the activity.



# INTACT

INDIGENOUS PEOPLES TRANSFORMING  
ALZHEIMER'S CARE TRAINING

## Questions and Answers



WORKING TOGETHER FOR HEALTH EQUITY



# IREACH

INSTITUTE FOR RESEARCH AND EDUCATION  
TO ADVANCE COMMUNITY HEALTH

